

Feeding Life Stages

LAYER

STARTING FROM HATCHING

This is a complete ration. Feed free choice from day of hatch until approximately 10 weeks of age. Provide fresh clean water at all times.

20%

<u>Nutrient</u>	<u>Chick Starter</u>
Crude Protein, min.....	20.00%
Lysine, min.....	1.00%
Methionine, min.....	0.40%
Crude Fat, min	3.00%
Crude Fiber, max.....	5.00%
Calcium(CA), min.....	.50%
Calcium (CA), max.....	1.00%
Phosphorus(P), min.....	0.60%
Salt (NaCl), min.....	0.20%
Salt (NaCl), max	0.60%

Approximate needs for 10 birds from Hatch to 10 weeks is 60 pounds.

GROWING OF PULLETS

Feed continuously as the sole ration from 10 weeks of age until birds begin laying at approximately 16 weeks of age. Provide fresh clean water at all times.

17%

<u>Nutrient</u>	<u>Pullet Grower</u>
Crude Protein, min.....	17.00%
Lysine, min.....	0.85%
Methionine, min.....	0.35%
Crude Fat, min	3.00%
Crude Fiber, max.....	5.00%
Calcium(CA), min.....	1.00%
Calcium (CA), max.....	1.50%
Phosphorus(P), min.....	0.70%
Salt (NaCl), min.....	0.30%
Salt (NaCl), max	0.70%

Approximate needs for 10 birds 10 weeks to laying is 120 pounds.

SUPPORTING LAYING HENS

17% Layer (Crumble/Pellet)

Feed as the complete ration for laying hens during periods of egg production. Feed intake will vary based upon size of birds and stage of laying. Fresh water should be available at all times. Additional grit and/or oyster shells should be provided.

Nutrient

Crude Protein, min.....	17.00%
Lysine, min.....	0.80%
Methionine, min.....	0.40%
Crude Fat, min.	3.50%
Crude Fiber, max.....	5.00%
Calcium(CA), min.....	3.90%
Calcium (CA), max.....	4.60%
Phosphorus(P), min.	0.70%
Salt (NaCl), min.....	0.25%
Salt (NaCl), max	0.70%

Approximate needs for 10 laying hens is 2.5 pounds per day.

20% Egg Mash

Prince 20% Egg Mash can be mixed with grain to make a complete egg ration or it can be fed free choice with scratch grain fed in a separate feeder or on the floor. Provide fresh clean water at all times.

Nutrient

Crude Protein, min.....	20.00%
Lysine, min.....	1.00%
Methionine, min.....	0.50%
Crude Fat, min.	3.50%
Crude Fiber, max.....	5.00%
Calcium (CA), min.....	3.25%
Calcium (CA), max.....	3.85%
Phosphorus(P), min.	0.70%
Salt (NaCl), min.....	0.30%
Salt (NaCl), max	0.70%

Approximate needs for 10 laying hens is 2.5 pounds per day.



Prince[®]
CORPORATION