

Feeding Life Stages

TURKEY

STARTING FROM HATCHING

Feed to newly arrived turkey pullets up to 10 weeks of age or up to slaughter.

Nutrient	28% Turkey Starter
Crude Protein (Min.)	28.00%
Lysine (Min.)	1.60%
Methionine (Min.)	0.60%
Crude Fat (Min.)	2.50%
Crude Fiber (Max.)	5.50%
Calcium (Min.)	1.00%
Calcium (Max.)	1.40%
Phosphorus (Min.)	0.95%
Salt (Min.)	0.35%
Salt (Max.)	0.60%

Feed as complete ration to newly hatched turkeys up to 10 weeks of age or up to slaughter. Provide fresh clean water at all times. Approximate need for 10 birds from Hatch to harvest is 500 pounds. (Assume 18 pound turkeys with even number of toms and hens) 28% Turkey Starter is available in a medicated form (Amprolium/Bacitracin) and a non-medicated from.

FINISHING

Feed as sole ration from 10 weeks of age until harvest.

Nutrient	18% Turkey Finisher
----------	---------------------

MEDICATED:

As an aid in prevention of coccidiosis.

ACTIVE DRUG INGREDIENTS:

Amprolium	0.0125% (113.5g/ton)
Crude Protein, min.	18.00%
Lysine, min.	1.00%
Methionine, min.	0.38%
Crude Fat, min.	3.00%
Crude Fiber, max.	5.50%
Calcium(CA), min.	0.90%
Calcium (CA), max.	1.20%
Phosphorus(P), min.	0.80%
Salt (NaCl), min.	0.35%
Salt (NaCl), max.	0.60%

Feed as a complete ration. Always provide plenty of clean water. Approximate need for 10 birds from 10 weeks to harvest is 350 pounds.



Prince[®]
CORPORATION