

Feeding Life Stages

BROILER

STARTING FROM HATCHING

Feed as the complete ration to newly hatched birds to 4 weeks of age or up to slaughter. Provide fresh clean water at all times.

25%

<u>Nutrient</u>	<u>Broiler Ration</u>
Crude Protein, min.....	25.00%
Lysine, min.....	1.40%
Methionine, min.....	0.50%
Crude Fat, min.	5.00%
Crude Fiber, max.....	5.00%
Calcium(CA), min.....	1.00%
Calcium (CA), max.....	1.50%
Phosphorus(P), min.....	0.90%
Salt (NaCl), min.....	0.40%
Salt (NaCl), max	0.80%

Approximate need for 10 birds from Hatch to harvest is 125 pounds.

25% Broiler Ration is available in a medicated form (Amprolium/Bacitracin) and a non-medicated from.



FINISHING BROILER

Feed as the complete ration to broilers from 4 weeks of age to finish. Provide fresh clean water at all times.

20%

<u>Nutrient</u>	<u>Broiler Finisher</u>
Crude Protein, min.....	20.00%
Lysine, min.....	1.18%
Methionine, min.....	0.50%
Crude Fat, min.	5.00%
Crude Fiber, max.....	5.00%
Calcium(CA), min.....	.80%
Calcium (CA), max.....	1.30%
Phosphorus(P), min.....	0.70%
Salt (NaCl), min.....	0.30%
Salt (NaCl), max	0.70%

Approximate need for 10 birds from 4 weeks to harvest is 80 pounds.

Prince[®]
CORPORATION